

Year 11 Math Homework

Student Name: _____	Grade: _____
Date: _____	Score: _____

Table of contents

4 Year 11 Topic 4 — Numbers and Functions (Part 1)	1
4.1 Numbers and Functions (Revision)	1
4.1.1 Surds and their Arithmetic	1
4.1.2 Equality of Surdic Expressions	3
4.1.3 Relations and Functions	4
4.1.4 Inverse Relations and Functions	6

This edition was printed on February 5, 2017 with worked solutions.
 Camera ready copy was prepared with the **L^AT_EX₂ ϵ** typesetting system.
 Copyright © 2000 - 2017 Yimin Math Centre

4 Year 11 Topic 4 — Numbers and Functions (Part 1)

4.1 Numbers and Functions (Revision)

4.1.1 Surds and their Arithmetic

Exercise 4.1.1

1. Use the result $\frac{\sqrt{a}}{\sqrt{b}} = \sqrt{\frac{a}{b}}$ to simplify these fractions:

(a) $\frac{\sqrt{72}}{\sqrt{98}}$ _____

(b) $\frac{\sqrt{50}}{\sqrt{8}}$ _____

(c) $\frac{\sqrt{52}}{\sqrt{91}}$ _____

(d) $\frac{\sqrt{175}}{\sqrt{28}}$ _____

2. Simplify each of these expressions completely:

(a) $\sqrt{96} - \sqrt{24} - \sqrt{54}$ _____

(b) $\sqrt{45} + \sqrt{80} - \sqrt{125}$ _____

(c) $\sqrt{63} + \sqrt{72} - \sqrt{50}$ _____

(d) $\sqrt{20} - \sqrt{12} + \sqrt{108}$ _____

3. Expand the following expressions and simplify them:

(a) $(\sqrt{3} - 1)(\sqrt{2} - 1)$ _____

(b) $(\sqrt{a} - 1)(\sqrt{a} + 1)$ _____

(c) $(2\sqrt{5} + \sqrt{3})(2 - \sqrt{3})$ _____

(d) $(\sqrt{x+1} + \sqrt{x-2})^2$

Exercise 4.1.2 Rationalising the denominator

1. Fully simplify these fractions:

(a) $\frac{6\sqrt{3} \times 5\sqrt{2}}{\sqrt{12} \times \sqrt{18}}$

(b) $\frac{5\sqrt{44} \times \sqrt{14}}{\sqrt{24} \times 3\sqrt{33}}$

2. Simplify the following by rationalising the denominator of each fraction:

(a) $\frac{1}{3+\sqrt{6}} - \frac{2}{\sqrt{6}}$

(b) $\frac{1}{3\sqrt{2}+1} + \frac{1}{1-3\sqrt{2}}$

3. Determine, without using a calculator, which is the greater number in each pair:

(a) $15 - 7\sqrt{2}$ or $3 + 2\sqrt{2}$

(b) $2\sqrt{6} - 3$ or $7 - 2\sqrt{6}$

4.1.2 Equality of Surdic Expressions**Exercise 4.1.3**

1. Find the value of integers x , y and z , given that z has no squares as factors:

(a) $x + y\sqrt{3} = (6 + \sqrt{3})^2$

(b) $x + y\sqrt{z} = (3 + \sqrt{5})^2$

2. Find the rational numbers a and b such that:

(a) $a + b\sqrt{3} = \frac{1}{2-\sqrt{3}}$

(b) $a + b\sqrt{6} = \frac{2\sqrt{6}+1}{2\sqrt{6}-3}$

3. Find the rational value of a and b , with $a > 0$ by forming two simultaneous equations and solving them: $(a + b\sqrt{2})^2 = 3 + 2\sqrt{2}$

4.1.3 Relations and Functions

- A function is a set of ordered pairs in which no two ordered pairs have the same x-coordinate.
- The domain of a function is the set of all x-coordinates of the ordered pairs.
- The range of a function is the set of all y-coordinates.

Exercise 4.1.4

1. Given that $f(x) = x^3 - x + 1$, evaluate and simplify the following:

(a) $\frac{f(h)-f(0)}{h}$

(b) $\frac{1}{6} (f(0) + 4f(\frac{1}{2}) + f(1))$

2. Find the natural domains of the following:

(a) $f(x) = \sqrt{9 - x^2}$

(b) $f(x) = \frac{1}{x^2 - 5x + 6}$

(c) $g(x) = \frac{x-3}{x^2-9}$

3. If $f(x) = \frac{1}{1-x}$, find $f(a-b)$.

Exercise 4.1.5

1. If $f(x) = 3^x$, show that $f(-x) = \frac{1}{f(x)}$

2. If $h(x) = \frac{x}{x^2-1}$, show that $h(\frac{1}{x}) = -h(x)$ for $x \neq 0$

3. If $f(x) = x + \frac{1}{x}$, show that $f(x) \times f(x + \frac{1}{x}) = f(x^2) + 3$

4. Given the functions $f(x) = x^2$, $F(x) = x - 3$, $g(x) = 2^x$ and $G(x) = 3x$, find:

(a) $F(f(x))$

(b) $G(g(x))$

4.1.4 Inverse Relations and Functions

- The inverse relation is obtained by reversing the values of each ordered pair.
- The domain of the inverse is the range of the relation and the range of the inverse is the domain of the relation.
- The graph of the inverse relation is obtained by reflecting the original graph in the line $y = x$.
- To find the equations and conditions of the inverse relation, write x for y , y for x and then solve for y .
- The inverse relation of a given relation is a function if and only if no horizontal line crosses the original graph more than once.

Exercise 4.1.6 Find the inverse algebraically by swapping x and y and then making y the subject:

1. $y = \frac{1}{x-1}$

2. $y = \frac{x+3}{x-3}$

3. $y = \frac{2x}{x+3}$

4. $y = \frac{2x-2}{x-2}$

Exercise 4.1.7

1. Each pair of functions $f(x)$ and $g(x)$ are mutually inverse. Verify in each case by substitution that:

(i) $f(g(2)) = 2$ and (ii) $g(f(2)) = 2$:

(a) $f(x) = x + 13$ and $g(x) = x - 13$

(b) $f(x) = x^3 - 6$ and $g(x) = \sqrt[3]{x + 6}$

2. Show that the inverse function of $y = \frac{ax+b}{x+c}$ is $y = \frac{b-cx}{x-a}$, for $x \leq 0$.

3. Hence show that $y = \frac{ax+b}{x+c}$ is its own inverse if and only if $a + c = 0$.

Exercise 4.1.8 Express $1 - \frac{2}{1-x}$ as a single fraction and hence find its reciprocal.

Exercise 4.1.9

1. If $f(x) = \frac{1}{1-x}$, find $f(f(x))$.

2. Find the range of the function $f(x) = \frac{1}{x^2+4x+7}$.

3. Solve $\frac{x}{1-x} > \frac{1}{3}$.

4. Solve $\frac{1}{1-x^2} \leq 4$.

5. Solve $|1 + 3x| = x - 2$.

Exercise 4.1.10

1. Show that $\frac{x^2+1}{x^2+4} = 1 - \frac{3}{x^2+4}$

2. Simplify $\frac{3^n+3^{n-2}}{3^{n-1}}$.

3. The equation $\frac{1}{1+x^2} = k$ has two distinct real roots. Find the possible values of k .

4. Prove that $\frac{1}{5-\sqrt{3}} + \frac{1}{5+\sqrt{3}}$ is rational.

5. Solve $x + |x| = 4$
